arXiv:1302.0501v1 [gr-qc] 3 Feb 2013

Many-body problem in Kaluza-Klein models with toroidal compactification

Alexey Chopovsky[1],* Maxim Eingorn[2,3],† and Alexander Zhuk[2]‡
[1] *Department of Theoretical Physics,*
Odessa National University,
Street Dvoryanskaya 2, 65082 Odessa, Ukraine

[2] *Astronomical Observatory, Odessa National University,*
Street Dvoryanskaya 2, 65082 Odessa, Ukraine
and
[3] *Department of Theoretical and Experimental Nuclear Physics,*
Odessa National Polytechnic University,
Shevchenko av. 1, 65044 Odessa, Ukraine

In this paper, we consider a system of gravitating bodies in Kaluza-Klein models with toroidal compactification of extra dimensions. To simulate the ordinary astrophysical objects (e.g., our Sun), we suppose that these bodies have nonrelativistic (pressureless) equations of state in the external/our space. At the same time, they may have nonzero parameters $\omega_{(\bar{\alpha}-3)}$ $(\bar{\alpha} = 4, \ldots, D)$ of the equations of state in the extra dimensions. We show that the presence of pressure/tension in the extra dimensions leads necessarily to the uniform smearing of the gravitating bodies over the internal space. Then, we prove that the Lagrange function of this many-body system can be constructed only if $\Sigma = \sum_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)} = 0$. This strongly contradicts the observations which demand $\Sigma \approx (3-D)/2 \sim O(1)$ for $D > 3$. Therefore, considered multidimensional Kaluza-Klein models face a severe problem.

PACS numbers: 04.25.Nx, 04.50.Cd, 04.80.Cc, 11.25.Mj

I. INTRODUCTION

The idea of multidimensionality of our Universe demanded by the theories of unification of the fundamental interactions is one of the most breathtaking ideas of theoretical physics. It takes its origin from the pioneering papers by Th. Kaluza and O. Klein [1], and now the most self-consistent modern theories of unification such as superstrings, supergravity and M-theory are constructed in spacetimes with extra dimensions (see, e.g., [2]). Different aspects of the idea of multidimensionality are intensively used in numerous modern articles.

Therefore, it is important to find experimental evidence for the existence of the extra dimensions. For example, one of the aims of Large Hadronic Collider consists in detecting of Kaluza-Klein (KK) particles which correspond to excitations of the internal spaces (see, e.g., [3]).

On the other hand, if we can show that the existence of the extra dimensions is contrary to observations, then these theories are prohibited.

In our previous papers [4–6] devoted to KK models with toroidal compactification of the extra dimensions, we have shown that gravitating masses should have tension in the internal space to be in agreement with gravitational experiments in the Solar system. For example, black strings/branes with the parameter $\omega = -1/2$ of the equation of state in the internal space satisfy this condition. For this value of ω, the variations of the internal space volume are absent [7]. In the dust-like case with $\omega = 0$, such variations generate the fifth force, that leads to contradictions with the experimental data. Hence, for individual gravitating bodies, we have shown in which cases and why the KK models with toroidal compactification are in agreement with the observations. Is it sufficient to claim that these models are viable?

To answer this question, in the present paper we continue the investigation of the KK models with toroidal compactification. Here, we try to construct the Lagrange function for a many-body system in these models. It is evident that any viable physical theory should permit such procedure. It is well known that the ordinary (nonrelativistic) objects, e.g., our Sun, have pressure that is much less than their energy density. Therefore, we suppose that the gravitating bodies for the considered system are pressureless in the external/our space. At the same time, we admit that they may have nonzero parameters $\omega_{(\bar{\alpha}-3)}$ $(\bar{\alpha} = 4, \ldots, D)$ of the equations of state in the extra dimensions.

To construct the Lagrange function for the many-body system, we should define preliminarily the gravitational field of this system. At this stage, our analysis leads to the first important conclusion: the assumption of the presence of pressure/tension in the extra dimensions requires necessarily the uniform smearing of the gravitating bodies over the internal space. Obviously, in the case of uniform smearing, the excited KK states are absent.

---

*Electronic address: alexey.chopovsky@gmail.com
†Electronic address: maxim.eingorn@gmail.com
‡Electronic address: ai.zhuk2@gmail.com

Then, we construct the Lagrange function of the considered many-body system and we come into conflict. On the one hand, the Lagrange function exists if the sum $\Sigma \equiv \sum_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)}$ of the parameters of the equations of state in the extra dimensions is equal to zero: $\Sigma = 0$. On the other hand, the agreement with the gravitational experiments requires $\Sigma \sim O(1)$. Thus, we claim that considered multidimensional KK models face a severe problem.

The paper is structured as follows. In Sec. II, we provide the general description of the model. In Sec. III and Sec. IV, we obtain the $1/c^2, 1/c^3$ and $1/c^4$ correction terms to the metric coefficients for the considered many-body system. In Sec. V, we demonstrate that gauge conditions lead to the uniform smearing of the gravitating bodies over the extra dimensions. The Lagrange function for the many-body system is constructed in the Sec. VI. The main results are summarized in concluding Sec. VII.

II. GENERAL SETUP

To construct the Lagrange function of a system of N massive bodies in $(D+1)$-dimensional spacetime, we define first the nonrelativistic gravitational field created by this system. To do it, we need to get the metric coefficients in the weak field limit. The general form of the multidimensional metrics is

$$ds^2 = g_{ik}dx^i dx^k = g_{00}\left(dx^0\right)^2 + 2g_{0\mu}dx^0 dx^\mu + g_{\mu\nu}dx^\mu dx^\nu \,, \tag{1}$$

where the Latin indices $i, k = 0, 1, \ldots, D$ and the Greek indices $\mu, \nu = 1, \ldots, D$. We make the natural assumption that in the case of the absence of matter sources the spacetime is Minkowski spacetime: $g_{00} = \eta_{00} = 1$, $g_{0\mu} = \eta_{0\mu} = 0$, $g_{\mu\nu} = \eta_{\mu\nu} = -\delta_{\mu\nu}$. In our paper, we consider in detail the case where the extra dimensions have the topology of tori. In the presence of matter, the metrics is not the Minkowskian one, and we investigate it in the weak field limit. It means that the gravitational field is weak and velocities of test bodies are small compared with the speed of light c. In the weak field limit the metrics is only slightly perturbed from its flat spacetime value. We will define the metrics (1) up to $1/c^2$ correction terms. Because the coordinate $x^0 = ct$, the metric coefficients can be expressed as follows:

$$\begin{aligned} &g_{00} \approx 1 + h_{00} + f_{00}\,, \\ &g_{0\mu} \approx h_{0\mu} + f_{0\mu}\,, \quad g_{\mu\nu} \approx -\delta_{\mu\nu} + h_{\mu\nu}\,, \end{aligned} \tag{2}$$

where $h_{ik} \sim O(1/c^2), f_{00} \sim O(1/c^4)$ and $f_{0\mu} \sim O(1/c^3)$. In particular, $h_{00} \equiv 2\varphi/c^2$. Later we will demonstrate that φ is the nonrelativistic gravitational potential. To get these correction terms, we should solve (in the corresponding orders of $1/c$) the multidimensional Einstein equation

$$R_{ik} = \frac{2S_D\tilde{G}_{\mathcal{D}}}{c^4}\left(T_{ik} - \frac{1}{D-1}g_{ik}T\right)\,, \tag{3}$$

where $S_D = 2\pi^{D/2}/\Gamma(D/2)$ is the total solid angle (the surface area of the $(D-1)$-dimensional sphere of the unit radius), $\tilde{G}_{\mathcal{D}}$ is the gravitational constant in the $(\mathcal{D} = D+1)$-dimensional spacetime. We consider a system of N discrete massive (with rest masses m_p, $p = 1, \ldots, N$) bodies. We suppose that the pressure of these bodies in the external three-dimensional space is much less than their energy density. This is a natural approximation for ordinary astrophysical objects such as our Sun. For example, in general relativity, this approach works well for calculating the gravitational experiments in the Solar system [8]. In other words, the gravitating bodies are pressureless in the external/our space. On the other hand, we suppose that they may have pressure in the extra dimensions. Therefore, nonzero components of the energy-momentum tensor of the system can be written in the following form:

$$T^{ik} = \tilde{\rho}c^2 u^i u^k, \quad i, k = 0, \ldots, 3\,, \tag{4}$$

$$T^{i\bar{\alpha}} = \tilde{\rho}c^2 u^i u^{\bar{\alpha}}\,, \quad i = 0, \ldots, 3;\ \bar{\alpha} = 4, \ldots, D\,, \tag{5}$$

$$T^{\bar{\alpha}\bar{\beta}} = -p_{(\bar{\alpha}-3)}g^{\bar{\alpha}\bar{\beta}} + \tilde{\rho}c^2 u^{\bar{\alpha}}u^{\bar{\beta}},\ \bar{\alpha}, \bar{\beta} = 4, \ldots, D \tag{6}$$

where the $(D+1)$-velocity $u^i = dx^i/ds$ and

$$\tilde{\rho} \equiv \sum_{p=1}^{N}\left[(-1)^D g\right]^{-1/2} m_p \sqrt{g_{lm}\frac{dx^l}{dx^0}\frac{dx^m}{dx^0}}\,\delta(\mathbf{x} - \mathbf{x}_p)\,, \tag{7}$$

where $\mathbf{x}_p$ is a D-dimensional radius-vector of the p-th particle. In what follows, the Greek indices $\alpha, \beta = 1, 2, 3$; $\bar{\alpha}, \bar{\beta} = 4, \ldots, D$ and μ, ν still run from 1 to D. In the extra dimensions we suppose the equations of state:

$$p_{(\bar{\alpha}-3)} = \omega_{(\bar{\alpha}-3)}\tilde{\rho}c^2\,. \tag{8}$$

If all parameters $\omega_{(\bar{\alpha}-3)} = 0$, then we come back to the model considered in our paper [4]. Here, massive bodies have dust-like equations of state in all spatial dimensions. If all $\omega_{(\bar{\alpha}-3)} = -1/2$ (tension in the extra dimensions), then these equations of state correspond to black strings (in the case of one extra dimension, i.e. $D = 4$) and black branes (for $D > 4$). If parameters satisfy the condition $\sum\limits_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)} \equiv \Sigma = -(D-3)/2$, then this case corresponds to latent solitons [6]. Obviously, black strings/branes satisfy this condition.

Now, we will solve the Einstein equation (3) in the corresponding orders of $1/c$. Obviously, for $\omega_{(\bar{\alpha}-3)} = 0\,, \bar{\alpha} = 4, \ldots, D$, we should reproduce the results of the paper [4].

III. $1/c^2$ CORRECTION TERMS

Here, we want to get the correction terms of the order $O(1/c^2)$. Since the right-hand side of Eq. (3) has the prefactor $1/c^4$, we should drop the terms of the order $O(c^n)$ with $n < 2$ in components of the energy-momentum tensor (4)-(6). Then, these components are approximated as

$$\begin{aligned}
T^{00} &\approx T_{00} \approx \rho c^2, \quad T^{\bar{\alpha}\bar{\beta}} \approx T_{\bar{\alpha}\bar{\beta}} \approx \omega_{(\bar{\alpha}-3)}\, \rho c^2\, \delta_{\bar{\alpha}\bar{\beta}}, \\
T^{0\mu} &\approx -T_{0\mu} \approx 0, \quad T^{\alpha\beta} \approx T_{\alpha\beta} \approx 0, \\
T &= T^{ik} g_{ik} \approx \rho c^2 (1-\Sigma)\,, \qquad (9)
\end{aligned}$$

where

$$\Sigma \equiv \sum_{\bar{\alpha}=4}^{D} \omega_{(\bar{\alpha}-3)} \qquad (10)$$

and we introduced the rest-mass density

$$\rho(\mathbf{x}) = \sum_{p=1}^{N} m_p \delta(\mathbf{x}-\mathbf{x}_p)\,. \qquad (11)$$

The left-hand side of (3) should also be of the order $O(1/c^2)$. Hence, the components of the Ricci tensor read [4]:

$$R_{00} \approx \frac{1}{2}\triangle_D h_{00}\,, \quad R_{0\mu} \approx \frac{1}{2}\triangle_D h_{0\mu}\,, \quad R_{\mu\nu} \approx \frac{1}{2}\triangle_D h_{\mu\nu}\,, \qquad (12)$$

where $\triangle_D = \delta^{\mu\nu}\partial^2/\partial x^\mu \partial x^\nu$ is the D-dimensional Laplace operator. Therefore, the Einstein equation (3) is reduced to the following system of equations:

$$\begin{aligned}
\triangle_D h_{00} &= \frac{4 S_D \tilde{G}_{\mathcal{D}}}{c^2} \frac{D-2+\Sigma}{D-1} \rho, \quad \triangle_D h_{0\mu} = 0\,, \\
\triangle_D h_{\alpha\beta} &= \frac{4 S_D \tilde{G}_{\mathcal{D}}}{c^2} \frac{D-2+\Sigma}{D-1} \frac{1-\Sigma}{D-2+\Sigma} \rho\, \delta_{\alpha\beta}\,, \\
\triangle_D h_{\bar{\alpha}\bar{\beta}} &= \frac{4 S_D \tilde{G}_{\mathcal{D}}}{c^2} \frac{\omega_{(\bar{\alpha}-3)}(D-1)+1-\Sigma}{D-1} \rho\, \delta_{\bar{\alpha}\bar{\beta}}\,,
\end{aligned} \qquad (13)$$

which have the following solutions

$$h_{00} = \frac{2\varphi(\mathbf{x})}{c^2}, \quad h_{0\mu} = 0, \qquad (14)$$

$$h_{\alpha\beta} = \frac{1-\Sigma}{D-2+\Sigma} \frac{2\varphi(\mathbf{x})}{c^2} \delta_{\alpha\beta}, \qquad (15)$$

$$h_{\bar{\alpha}\bar{\beta}} = \frac{\omega_{(\bar{\alpha}-3)}(D-1)+1-\Sigma}{D-2+\Sigma} \frac{2\varphi(\mathbf{x})}{c^2} \delta_{\bar{\alpha}\bar{\beta}}\,, \qquad (16)$$

where the function $\varphi(\mathbf{x})$ satisfies the D-dimensional Poisson equation

$$\triangle_D \varphi(\mathbf{x}) = 2 S_D \tilde{G}_{\mathcal{D}} \frac{D-2+\Sigma}{D-1} \rho(\mathbf{x})\,. \qquad (17)$$

We would remind that $\mathbf{x}$ is a D-dimensional radius-vector. It is worth noting that if $\omega_{(\bar{\alpha}-3)} = 0,\ \forall\ \bar{\alpha} \ \Rightarrow \Sigma = 0$, then we reproduce the results of the paper [4]. On the other hand, if all $\omega_{(\bar{\alpha}-3)} = -1/2$, then $h_{\bar{\alpha}\bar{\beta}} = 0$ that should take place for black strings/branes [6].

IV. $1/c^3$ AND $1/c^4$ CORRECTION TERMS

Now, to find the corrections f_{00} and $f_{0\mu}$ in (2), we should take into account also the terms of the orders $O(c^0)$ and $O(c^1)$ in T_{00} and $T_{0\mu}$, respectively. It can be easily seen that the components $T_{\alpha\beta}$ and $T_{\alpha\bar{\beta}}$ are of the order of $O(c^0)$ and, due to the prefactor $1/c^4$ in the right-hand side of (3), they can contribute to $g_{\alpha\beta}$ and $g_{\alpha\bar{\beta}}$ terms of the order $O(1/c^4)$ which is not of interest for us. Therefore, we will take these components into account just to calculate the trace $T = g_{ik}T^{ik}$ where we will keep the terms $O(c^2)$ and $O(c^0)$. Concerning the components $T_{\bar{\alpha}\bar{\beta}}$, besides the terms $O(c^2)$, we should also calculate the terms $O(c^0)$. Additionally, within our accuracy we can conclude that

$$\begin{aligned}
T_{00} &\approx \left(1+\frac{4\varphi}{c^2}\right) T^{00}, \\
T_{0\mu} &\approx -T^{0\mu}, \quad T_{\alpha\beta} \approx T^{\alpha\beta}, \quad T_{\alpha\bar{\beta}} \approx T^{\alpha\bar{\beta}}\,, \qquad (18) \\
T_{\bar{\alpha}\bar{\beta}} &\approx \left(1-\frac{4\varphi}{c^2}\frac{\omega_{(\bar{\alpha}-3)}(D-1)+1-\Sigma}{D-2+\Sigma}\right) T^{\bar{\alpha}\bar{\beta}}\,.
\end{aligned}$$

To calculate these covariant components, we will also use the following auxiliary expressions:

$$\sqrt{g_{lm}\frac{dx^l}{dx^0}\frac{dx^m}{dx^0}} \approx 1 + \frac{\varphi - v^2/2}{c^2},$$

$$[(-1)^D g]^{-1/2} \approx 1 + \frac{2\varphi}{c^2}\frac{1-\Sigma}{D-2+\Sigma},$$

$$\tilde{\rho} \approx \rho \left[1 + \frac{\varphi}{c^2}\frac{D-\Sigma}{D-2+\Sigma} - \frac{v^2}{2c^2}\right]\,,$$

$$u^0 \approx 1 - \frac{\varphi - v^2/2}{c^2}, \quad u^\mu \approx \frac{v^\mu}{c},$$

where $v^\mu = dx^\mu/dt$ and $v^2 = \delta_{\mu\nu}v^\mu v^\nu$.

Taking into account all comments made above, we get the covariant components of the energy-momentum tensor

$$T_{00} \approx \rho c^2 \left[1 + \frac{\varphi}{c^2}\frac{3D-4+\Sigma}{D-2+\Sigma} + \frac{v^2}{2c^2}\right]\,, \qquad (19)$$

$$T_{0\mu} \approx -\rho c v^\mu\,, \qquad (20)$$

$$T_{\alpha\beta} \approx \rho v^\alpha v^\beta\,, \quad T_{\alpha\bar{\beta}} \approx \rho v^\alpha v^{\bar{\beta}}\,, \qquad (21)$$

$$\begin{aligned}
T_{\bar{\alpha}\bar{\beta}} &\approx \rho c^2 \left\{\omega_{(\bar{\alpha}-3)}\delta_{\bar{\alpha}\bar{\beta}}\right. \\
&\times \left[1 + \frac{\varphi}{c^2}\frac{D-\Sigma-2\left[\omega_{(\bar{\alpha}-3)}(D-1)+1-\Sigma\right]}{D-2+\Sigma} - \frac{v^2}{2c^2}\right] \\
&+ \left.\frac{v^{\bar{\alpha}}v^{\bar{\beta}}}{c^2}\right\} \qquad (22)
\end{aligned}$$

and the trace

$$T \approx \rho c^2(1-\Sigma) + \rho\varphi\frac{(D-\Sigma)(1-\Sigma)}{D-2+\Sigma} + \rho(\Sigma-1)\frac{v^2}{2}\,. \quad (23)$$

For the 00 and 0μ components of the Ricci tensor we get (see the details in [4]):

$$\begin{aligned} R_{00} &\approx \frac{1}{c^2}\triangle_D\varphi + \frac{1}{2}\triangle_D\left(f_{00} - \frac{2}{c^4}\varphi^2\right) + \frac{2}{c^4}\frac{D-1}{D-2}\varphi\triangle_D\varphi, \\ R_{0\mu} &\approx \frac{1}{2}\triangle_D f_{0\mu} + \frac{1}{2c^3}\frac{\partial^2\varphi}{\partial t\partial x^\mu}\,. \end{aligned} \quad (24)$$

Then, the 00 component of the Einstein equation (3) reads

$$\begin{aligned} &\triangle_D\left(f_{00} - \frac{2}{c^4}\varphi^2\right) = \frac{4S_D\tilde{G}_{\mathcal{D}}}{c^4}\sum_p m_p \\ &\times \left[\frac{(D-2)^2 - \Sigma D}{(D-1)(D-2)}\varphi'(\mathbf{x}_p) + \frac{D-\Sigma}{2(D-1)}v_p^2\right]\delta(\mathbf{x}-\mathbf{x}_p)\,, \end{aligned} \quad (25)$$

where we used Eqs. (11) and (17). Here, v_p is the three-velocity of the p-th particle and $\varphi'(\mathbf{x}_p)$ is the potential of the gravitational field at a point with the radius vector $\mathbf{x}_p$ produced by all particles, except the p-th. Subtraction of the infinite contribution of the gravitational field of the p-th particle corresponds to the renormalization of its mass (see [8]). Obviously, if $\varphi_p(\mathbf{x}-\mathbf{x}_p)$ is the potential of the p-th particle (with the radius vector $\mathbf{x}_p$) at a point $\mathbf{x}$:

$$\triangle_D\varphi_p(\mathbf{x}-\mathbf{x}_p) = 2S_D\tilde{G}_{\mathcal{D}}\frac{D-2+\Sigma}{D-1}m_p\delta(\mathbf{x}-\mathbf{x}_p)\,, \quad (26)$$

then

$$\begin{aligned} \varphi(\mathbf{x}) &= \sum_p \varphi_p(\mathbf{x}-\mathbf{x}_p)\,, & (27) \\ \varphi'(\mathbf{x}_p) &= \sum_{q\neq p}\varphi_q(\mathbf{x}_p-\mathbf{x}_q)\,. & (28) \end{aligned}$$

The solution of Eq. (25) is

$$\begin{aligned} f_{00}(\mathbf{x}) &= \frac{2}{c^4}\varphi^2(\mathbf{x}) \\ &+ \frac{2}{c^4}\frac{(D-2)^2 - \Sigma D}{(D-2+\Sigma)(D-2)}\sum_p\varphi_p(\mathbf{x}-\mathbf{x}_p)\varphi'(\mathbf{x}_p) \\ &+ \frac{1}{c^4}\frac{D-\Sigma}{D-2+\Sigma}\sum_p\varphi_p(\mathbf{x}-\mathbf{x}_p)v_p^2\,. \end{aligned} \quad (29)$$

For the 0μ component of the Einstein equation we have

$$\begin{aligned} &\frac{1}{2}\triangle_D f_{0\mu} + \frac{1}{2c^3}\frac{\partial^2\varphi}{\partial t\partial x^\mu} = -\frac{2S_D\tilde{G}_{\mathcal{D}}}{c^3}\sum_p m_p v_p^\mu\delta(\mathbf{x}-\mathbf{x}_p) \\ &= -\frac{1}{c^3}\frac{D-1}{D-2+\Sigma}\sum_p\triangle_D\varphi_p(\mathbf{x}-\mathbf{x}_p)v_p^\mu\,, \end{aligned} \quad (30)$$

and the solution is

$$f_{0\mu}(\mathbf{x}) = -\frac{2}{c^3}\frac{D-1}{D-2+\Sigma}\sum_p\varphi_p(\mathbf{x}-\mathbf{x}_p)v_p^\mu - \frac{1}{c^3}\frac{\partial^2 f}{\partial t\partial x^\mu}, \quad (31)$$

where the function f satisfies the following equation:

$$\triangle_D f = \varphi(\mathbf{x})\,. \quad (32)$$

It can be easily verified that solutions (29) and (31) are reduced to the results of paper [4] for $\Sigma = 0$.

V. GAUGE CONDITIONS AND SMEARING

To get the expressions (12) and (24), we used the standard (see, e.g., Eq. (105.10) in [8]) gauge condition

$$\partial_k\left(h_i^k - \frac{1}{2}h_l^l\delta_i^k\right) = 0\,, \quad i,k = 0,1,\ldots,D\,, \quad (33)$$

where $h_i^k \equiv \eta^{km}h_{mi}$. Hence,

$$h_0^0 = \eta^{00}h_{00} = h_{00}, \quad h_\nu^\mu = \eta^{\mu\kappa}h_{\nu\kappa} = -h_{\mu\nu}\,. \quad (34)$$

Therefore,

$$\begin{aligned} h_0^0 &= \frac{2\varphi(\mathbf{x})}{c^2}, \quad h_\alpha^\beta = -\frac{1-\Sigma}{D-2+\Sigma}\,\frac{2\varphi(\mathbf{x})}{c^2}\delta_\alpha^\beta, & (35) \\ h_{\bar\alpha}^{\bar\beta} &= -\frac{\omega_{(\bar\alpha-3)}(D-1) + 1 - \Sigma}{D-2+\Sigma}\,\frac{2\varphi(\mathbf{x})}{c^2}\delta_{\bar\alpha}^{\bar\beta}\,, & (36) \\ h_l^l &= \frac{2(\Sigma-1)}{D-2+\Sigma}\,\frac{2\varphi(\mathbf{x})}{c^2}\,. & (37) \end{aligned}$$

Let us check that these solutions satisfy the condition (33). For $i = 0$, we get immediately

$$\partial_k\left(h_0^k - \frac{1}{2}h_l^l\delta_0^k\right) = \partial_0\left(h_0^0 - \frac{1}{2}h_l^l\right) = 0 + O\left(\frac{1}{c^3}\right)\,. \quad (38)$$

For $i = \beta$ we have

$$\begin{aligned} &\partial_k\left(h_\beta^k - \frac{1}{2}h_l^l\delta_\beta^k\right) = \partial_\alpha\left(h_\beta^\alpha - \frac{1}{2}h_l^l\delta_\beta^\alpha\right) \\ &= \left[-\frac{1-\Sigma}{D-2+\Sigma} + \frac{1-\Sigma}{D-2+\Sigma}\right]\frac{2}{c^2}\partial_\beta\varphi = 0\,, \end{aligned} \quad (39)$$

that is the condition is automatically satisfied. For $i = \bar\beta$ we obtain

$$\begin{aligned} &\partial_k\left(h_{\bar\beta}^k - \frac{1}{2}h_l^l\delta_{\bar\beta}^k\right) = \partial_{\bar\alpha}\left(h_{\bar\beta}^{\bar\alpha} - \frac{1}{2}h_l^l\delta_{\bar\beta}^{\bar\alpha}\right) \\ &= -\frac{\omega_{(\bar\beta-3)}(D-1)}{D-2+\Sigma}\frac{2}{c^2}\partial_{\bar\beta}\varphi = 0\,. \end{aligned} \quad (40)$$

To satisfy this condition, we should demand either $\omega_{(\bar\beta-3)} = 0$ or $\partial_{\bar\beta}\varphi = 0$. Because we consider the general case $\omega_{(\bar\beta-3)} \neq 0$, we must choose the latter condition. Therefore, the presence of nonzero pressure/tension in the extra dimensions results in the metric coefficients which do not depend on the coordinates of the internal space, i.e. the gravitating masses should be uniformly smeared over the extra dimensions. In this case, the rest mass density (11) should be rewritten in the form: $\rho(\mathbf{x}) \to \rho(\mathbf{r}) = \sum_p m_p \delta(\mathbf{r}-\mathbf{r}_p)/\prod_{\bar\alpha} a_{(\bar\alpha-3)}$, where $\mathbf{r}_p$ is a three-dimensional radius vector of the p-th particle in the external space, $a_{(\bar\alpha-3)}$ are periods of the tori (i.e. $\prod_{\bar\alpha} a_{(\bar\alpha-3)}$ is the volume of the internal space). Then, Eq. (17) is reduced to the ordinary three-dimensional Poisson equation

$$\triangle_3 \varphi(\mathbf{r}) = 4\pi G_N \sum_p m_p \delta(\mathbf{r}-\mathbf{r}_p) \tag{41}$$

with the solution

$$\varphi(\mathbf{r}) = -\sum_p \frac{G_N m_p}{|\mathbf{r}-\mathbf{r}_p|} = \sum_p \varphi_p(\mathbf{r}-\mathbf{r}_p)\,, \tag{42}$$

where G_N is the Newtonian gravitational constant:

$$4\pi G_N = \frac{2S_D(D-2+\Sigma)}{(D-1)\prod_{\bar\alpha} a_{(\bar\alpha-3)}}\tilde G_{\mathcal{D}}\,. \tag{43}$$

Hereafter, $\mathbf{r}, \mathbf{r}_p$ are radius vectors in three-dimensional external/our space.

In the case of the smearing, Eq. (32) has the following solution

$$f(\mathbf{r}) = -\frac{G_N}{2}\sum_p m_p |\mathbf{r}-\mathbf{r}_p|\,, \tag{44}$$

where, to get it, we used the well known equation $\triangle_3 r = 2/r$ in the three-dimensional flat space. Because

$$\frac{\partial}{\partial t}\left(\frac{\partial|\mathbf{r}-\mathbf{r}_p|}{\partial x^\alpha}\right) = \frac{\partial}{\partial t}\left(\frac{x^\alpha - x^\alpha_p}{|\mathbf{r}-\mathbf{r}_p|}\right) \tag{45}$$
$$= \frac{1}{|\mathbf{r}-\mathbf{r}_p|^2}\left[-v^\alpha_p|\mathbf{r}-\mathbf{r}_p| - \frac{x^\alpha - x^\alpha_p}{|\mathbf{r}-\mathbf{r}_p|}\sum_\beta (x^\beta - x^\beta_p)(-v^\beta_p)\right],$$

we get for $f_{0\alpha}$:

$$f_{0\alpha} = \frac{G_N}{2c^3}\sum_p \frac{m_p}{|\mathbf{r}-\mathbf{r}_p|}\left(\frac{3D-2-\Sigma}{D-2+\Sigma}v^\alpha_p + n^\alpha_p(\mathbf{n}_p\mathbf{v}_p)\right)\,, \tag{46}$$

where we introduce the three-dimensional unit vector in the direction from the p-th particle to a point with the radius vector $\mathbf{r}$:

$$n^\alpha_p = \frac{x^\alpha - x^\alpha_p}{|\mathbf{r}-\mathbf{r}_p|}, \tag{47}$$

and $(\mathbf{n}_p\mathbf{v}_p) = \sum_\beta n^\beta_p v^\beta_p$.

It should be noted that, to get the formula (31), we used the following gauge condition:

$$\frac{\partial f^\mu_0}{\partial x^\mu} - \frac{1}{2}\frac{\partial h^\mu_\mu}{\partial x^0} = 0\,, \tag{48}$$

where $f^\mu_0 = \eta^{k\mu} f_{0k} = -f_{0\mu}$. In the case of smearing, this condition is reduced to

$$\frac{\partial f^\beta_0}{\partial x^\beta} - \frac{1}{2}\frac{\partial h^\mu_\mu}{\partial x^0} = 0\,, \tag{49}$$

where we remind that $\alpha, \beta = 1,2,3$ and $\mu, \nu = 1, \ldots, D$. Taking into account the following auxiliary equations:

$$\frac{\partial}{\partial x^\beta}\frac{1}{|\mathbf{r}-\mathbf{r}_p|} = -\frac{n^\beta_p}{|\mathbf{r}-\mathbf{r}_p|^2}\,, \tag{50}$$

$$\sum_\beta \frac{\partial}{\partial x^\beta}\left(n^\beta_p \frac{(\mathbf{n}_p\mathbf{v}_p)}{|\mathbf{r}-\mathbf{r}_p|}\right) = \frac{(\mathbf{n}_p\mathbf{v}_p)}{|\mathbf{r}-\mathbf{r}_p|^2}\,, \tag{51}$$

$$\frac{\partial}{\partial t}\frac{1}{|\mathbf{r}-\mathbf{r}_p|} = \frac{(\mathbf{n}_p\mathbf{v}_p)}{|\mathbf{r}-\mathbf{r}_p|^2}\,, \tag{52}$$

we can easily seen that the condition (49) is satisfied:

$$\frac{\partial f^\beta_0}{\partial x^\beta} - \frac{1}{2}\frac{\partial h^\mu_\mu}{\partial x^0} = \frac{G_N}{2c^3}\left\{\frac{3D-2-\Sigma}{D-2+\Sigma}\sum_p m_p \frac{(\mathbf{n}_p\mathbf{v}_p)}{|\mathbf{r}-\mathbf{r}_p|^2}\right.$$
$$\left. - \sum_p m_p \frac{(\mathbf{n}_p\mathbf{v}_p)}{|\mathbf{r}-\mathbf{r}_p|^2} - \frac{2(D-\Sigma)}{D-2+\Sigma}\sum_p m_p\frac{(\mathbf{n}_p\mathbf{v}_p)}{|\mathbf{r}-\mathbf{r}_p|^2}\right\} = 0\,. \tag{53}$$

Because the presence of pressure/tension in the extra dimensions requires the uniform smearing of the gravitating masses over the internal space, we provide the metric coefficients in this case:

$$g_{00} \approx 1 + \frac{2\varphi(\mathbf{r})}{c^2} + \frac{2\varphi^2(\mathbf{r})}{c^4} + \frac{(D-2)^2 - \Sigma D}{(D-2+\Sigma)(D-2)}\frac{2G^2_N}{c^4}$$
$$\times \sum_p \frac{m_p}{|\mathbf{r}-\mathbf{r}_p|}\sum_{q\neq p}\frac{m_q}{|\mathbf{r}_p-\mathbf{r}_q|}$$
$$- \frac{D-\Sigma}{D-2+\Sigma}\frac{G_N}{c^4}\sum_p \frac{m_p v^2_p}{|\mathbf{r}-\mathbf{r}_p|}\,, \tag{54}$$

$$g_{0\alpha} \approx \frac{3D-2-\Sigma}{D-2+\Sigma}\frac{G_N}{2c^3}\sum_p \frac{m_p}{|\mathbf{r}-\mathbf{r}_p|}v^\alpha_p$$
$$+ \frac{G_N}{2c^3}\sum_p \frac{m_p}{|\mathbf{r}-\mathbf{r}_p|}n^\alpha_p(\mathbf{n}_p\mathbf{v}_p)\,, \tag{55}$$

$$g_{\alpha\beta} \approx \left(-1 + \frac{1-\Sigma}{D-2+\Sigma}\frac{2\varphi(\mathbf{r})}{c^2}\right)\delta_{\alpha\beta}\,, \tag{56}$$

$$g_{\bar\alpha\bar\beta} \approx \left(-1 + \frac{\omega_{(\bar\alpha-3)}(D-1)+1-\Sigma}{D-2+\Sigma}\frac{2\varphi(\mathbf{r})}{c^2}\right)\delta_{\bar\alpha\bar\beta}\,, \tag{57}$$

where the potential $\varphi(\mathbf{r})$ is given by (42).

VI. LAGRANGE FUNCTION FOR A MANY-BODY SYSTEM

Let us construct now the Lagrange function of the many-body system described above. To perform it, we will follow the procedure described in [8] (see §106). The Lagrange function of a particle a with the mass m_a in the gravitational field created by the other bodies is given by the expression

$$\begin{aligned} L_a &= -m_a c \frac{ds_a}{dt} \qquad (58)\\ &= -m_a c^2 \left(g_{00} + 2\sum_{\mu} g_{0\mu}\frac{v_a^{\mu}}{c} + \sum_{\mu\nu} g_{\mu\nu}\frac{v_a^{\mu}v_a^{\nu}}{c^2}\right)^{1/2}, \end{aligned}$$

where the metric coefficients are taken at $\mathbf{r} = \mathbf{r}_a$. We should keep in mind that in the case of the smeared (over the extra dimensions) gravitating masses, the components of the velocity in the extra dimensions are equal to zero: $v_a^{\mu} = (v_a^{\alpha}, v_a^{\bar{\alpha}}) = (v_a^{\alpha}, 0)$. It is convenient to rewrite the metric coefficients (54)-(56) in the following form:

$$\begin{aligned} g_{00} &\approx 1 + \frac{1}{c^2}\gamma_{00}^{(1)} + \frac{1}{c^4}\gamma_{00}^{(2)}, \quad g_{0\alpha} \approx \frac{1}{c^3}\gamma_{0\alpha}\,,\\ g_{\alpha\beta} &\approx \left(-1 + \frac{1}{c^2}\gamma_{(\alpha)}\right)\delta_{\alpha\beta}\,, \end{aligned}$$

where the meaning of the functions γ is evident. Then, we get

$$\begin{aligned} \frac{ds_a}{dt} &\approx c\left\{1 + \frac{1}{2c^2}\left[\gamma_{00}^{(1)} - v_a^2\right]\right.\\ &+ \frac{1}{2c^4}\left[\gamma_{00}^{(2)} + 2\sum_{\alpha}\gamma_{0\alpha}v_a^{\alpha} + \sum_{\alpha\beta}\gamma_{(\alpha)}\delta_{\alpha\beta}v_a^{\alpha}v_a^{\beta}\right]\\ &- \left.\frac{1}{8c^4}\left[\gamma_{00}^{(1)} - v_a^2\right]^2\right\}. \qquad (59) \end{aligned}$$

Substituting the explicit form of the metric coefficients (54)-(56), we obtain

$$\begin{aligned} L_a &= -m_a c^2 + \frac{m_a v_a^2}{2} + \frac{m_a v_a^4}{8c^2} + G_N\sum_{p}\frac{m_a m_p}{|\mathbf{r}-\mathbf{r}_p|}\\ &- \frac{1}{2c^2}G_N^2\sum_{p}\sum_{q}\frac{m_a m_p m_q}{|\mathbf{r}-\mathbf{r}_p||\mathbf{r}-\mathbf{r}_q|}\\ &- \frac{\mathfrak{a}(D,\Sigma)}{c^2}G_N^2\sum_{p}\sum_{q\neq p}\frac{m_a m_p m_q}{|\mathbf{r}-\mathbf{r}_p||\mathbf{r}_p-\mathbf{r}_q|}\\ &+ \frac{1}{2c^2}G_N\sum_{p}\frac{m_a m_p}{|\mathbf{r}-\mathbf{r}_p|}\left[\mathfrak{b}(D,\Sigma)v_p^2 + 2\left(\mathfrak{d}(D,\Sigma)+\frac{1}{2}\right)v_a^2\right.\\ &- \left.\mathfrak{c}(D,\Sigma)(\mathbf{v}_p\mathbf{v}_a) - (\mathbf{n}_p\mathbf{v}_p)(\mathbf{n}_p\mathbf{v}_a)\right]. \qquad (60) \end{aligned}$$

Here, we use the following abbreviations:

$$\begin{aligned} \mathfrak{a}(D,\Sigma) &\equiv \frac{(D-2)^2 - \Sigma D}{(D-2+\Sigma)(D-2)}, \quad \mathfrak{b}(D,\Sigma) \equiv \frac{D-\Sigma}{D-2+\Sigma},\\ \mathfrak{c}(D,\Sigma) &\equiv \frac{3D-2-\Sigma}{D-2+\Sigma}, \quad \mathfrak{d}(D,\Sigma) \equiv \frac{1-\Sigma}{D-2+\Sigma}. \qquad (61) \end{aligned}$$

We remind that in the expression (60) $\mathbf{r} = \mathbf{r}_a$ and all infinite terms should be cast out. For our purposes, it is sufficient to consider the case of two particles. Then, for the particle "1", we have the following expression:

$$\begin{aligned} L_1 &= \mathrm{f}(\mathbf{v}_1^2) + G_N\frac{m_1 m_2}{|\mathbf{r}-\mathbf{r}_2|} - \frac{1}{2c^2}G_N^2\frac{m_1 m_2^2}{|\mathbf{r}-\mathbf{r}_2|^2}\\ &- \frac{1}{c^2}\mathfrak{a}(D,\Sigma)G_N^2\frac{m_1^2 m_2}{|\mathbf{r}-\mathbf{r}_2||\mathbf{r}_1-\mathbf{r}_2|}\\ &+ \frac{1}{2c^2}\frac{G_N m_1 m_2}{|\mathbf{r}-\mathbf{r}_2|}\left[\mathfrak{b}(D,\Sigma)v_2^2 + 2\left(\mathfrak{d}(D,\Sigma)+\frac{1}{2}\right)v_1^2\right.\\ &- \left.\mathfrak{c}(D,\Sigma)(\mathbf{v}_1\mathbf{v}_2) - (\mathbf{n}_2\mathbf{v}_2)(\mathbf{n}_2\mathbf{v}_1)\right], \qquad (62) \end{aligned}$$

where $\mathrm{f}(\mathbf{v}_1^2) = m_1 v_1^2/2 + m_1 v_1^4/(8c^2)$ and we drop the term $-m_1 c^2$.

The total Lagrange function of the two-body system should be constructed so that it leads to the correct values of the forces $\partial L_a/\partial\mathbf{r}|_{\mathbf{r}=\mathbf{r}_a}$ acting on each of the bodies for given motion of the others [8]. To achieve it, we, first, will differentiate L_1 with respect to $\mathbf{r}$, setting $\mathbf{r} = \mathbf{r}_1$ after that. Then, we should integrate this expression with respect to $\mathbf{r}_1$. Following this prescription and taking into account a useful auxiliary relation

$$\begin{aligned} &\left(-\frac{1}{2c^2}G_N^2 m_1 m_2^2\frac{\partial}{\partial\mathbf{r}}\frac{1}{|\mathbf{r}-\mathbf{r}_2|^2}\right.\\ &- \left.\left.\frac{\mathfrak{a}(D,\Sigma)G_N^2}{c^2}\frac{m_1^2 m_2}{|\mathbf{r}_1-\mathbf{r}_2|}\frac{\partial}{\partial\mathbf{r}}\frac{1}{|\mathbf{r}-\mathbf{r}_2|}\right)\right|_{\mathbf{r}=\mathbf{r}_1}\\ &= -\frac{1}{2c^2}G_N^2 m_1 m_2(\mathfrak{a}m_1 + m_2)\frac{\partial}{\partial\mathbf{r}_1}\frac{1}{|\mathbf{r}_1-\mathbf{r}_2|^2}, \qquad (63) \end{aligned}$$

we obtain from (62) the two-body Lagrange function

$$\begin{aligned} L_1^{(2)} &= \tilde{\mathrm{f}}(\mathbf{v}_1^2,\mathbf{v}_2^2) + \frac{G_N m_1 m_2}{r_{12}} - \frac{G_N^2 m_1 m_2(\mathfrak{a}m_1 + m_2)}{2c^2 r_{12}^2}\\ &+ \frac{G_N m_1 m_2}{2c^2 r_{12}}\left[\mathfrak{b}(D,\Sigma)v_2^2 + (2\mathfrak{d}(D,\Sigma)+1)v_1^2\right.\\ &- \left.\mathfrak{c}(D,\Sigma)(\mathbf{v}_1\mathbf{v}_2) - (\mathbf{n}_{12}\mathbf{v}_1)(\mathbf{n}_{12}\mathbf{v}_2)\right], \qquad (64) \end{aligned}$$

where $\tilde{\mathrm{f}}(\mathbf{v}_1^2,\mathbf{v}_2^2) = \sum_{a=1}^{2} m_a v_a^2/2 + \sum_{a=1}^{2} m_a v_a^4/(8c^2)$ and $r_{12} \equiv |\mathbf{r}_1 - \mathbf{r}_2|$. It can be easily seen that $\partial L_1/\partial\mathbf{r}|_{\mathbf{r}=\mathbf{r}_1} = \partial L_1^{(2)}/\partial\mathbf{r}_1$. By the same way we can construct the two-body Lagrange function $L_2^{(2)}$ from the Lagrange function L_2 for the particle "2":

$$\begin{aligned} L_2^{(2)} &= \tilde{\mathrm{f}}(\mathbf{v}_1^2,\mathbf{v}_2^2) + \frac{G_N m_1 m_2}{r_{12}} - \frac{G_N^2 m_1 m_2(m_1 + \mathfrak{a}m_2)}{2c^2 r_{12}^2}\\ &+ \frac{G_N m_1 m_2}{2c^2 r_{12}}\left[\mathfrak{b}(D,\Sigma)v_1^2 + (2\mathfrak{d}(D,\Sigma)+1)v_2^2\right.\\ &- \left.\mathfrak{c}(D,\Sigma)(\mathbf{v}_1\mathbf{v}_2) - (\mathbf{n}_{12}\mathbf{v}_1)(\mathbf{n}_{12}\mathbf{v}_2)\right]. \qquad (65) \end{aligned}$$

It is worth noting that both $L_1^{(2)}$ and $L_2^{(2)}$ are reduced to the Lagrange function of the two-body system in [8] if we assume that $D = 3$, $\Sigma = 0$.

Obviously, the Lagrange functions $L_1^{(2)}$ and $L_2^{(2)}$ should be symmetric with respect to permutations of particles 1 and 2 and should coincide with each other. This requires the following conditions:

$$\mathfrak{a}(D, \Sigma) = 1, \quad \mathfrak{b}(D, \Sigma) = 2\mathfrak{d}(D, \Sigma) + 1 . \tag{66}$$

The second condition is satisfied identically. However, the former one is valid only for

$$\Sigma = 0 . \tag{67}$$

This happens, e.g., for the dust-like equations of state in the extra dimensions: $\omega_{(\bar{\alpha}-3)} = 0$, $\forall\, \bar{\alpha}$. Definitely, black strings/branes contradict this condition.

We can also get an additional restriction on Σ from the known gravitational experiments in the Solar system. For example, the post-Newtonian parameter γ for the considered models is expressed via Σ as follows (see, e.g., [4–6]):

$$\gamma - 1 = \frac{3 - D - 2\Sigma}{D - 2 + \Sigma} . \tag{68}$$

There are strong experimental restrictions on the value of γ. The tightest constraint on γ comes from the Shapiro time-delay experiment using the Cassini spacecraft, namely: $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ [9–11]. Therefore, $\gamma - 1 \approx 0 \;\Rightarrow\; \Sigma \approx (3 - D)/2 \sim O(1)$ for $D > 3$. It is worth noting that for latent solitons (and black strings/branes as their particular cases) $\Sigma = (3 - D)/2$ [6] and $\gamma = 1$, as in general relativity. This is in good agreement with observations.

Hence, we come into conflict. On the one hand, the Lagrange function for a two-body system exists for $\Sigma = 0$. On the other hand, this value contradicts the gravitational experiments.

VII. CONCLUSION

In this paper, we have constructed the Lagrange function for a two-body system in the case of Kaluza-Klein models with toroidal compactification of the extra dimensions. The case of more than two bodies is straightforward. We supposed that gravitating bodies are pressureless in the external/our space. This is a natural approximation for ordinary/nonrelativistic astrophysical objects such as our Sun, since the pressure in these bodies is much less than their energy density. For example, this approach works well for calculating the gravitational experiments in the Solar system [8]. On the other hand, we have seen from our previous investigations [5–7] that, to achieve the agreement with the gravitational experiments in KK models with toroidal compactification, the gravitating bodies should have pressure/tension in the extra dimensions. Therefore, in the present paper we supposed that gravitating masses may have nonzero parameters $\omega_{(\bar{\alpha}-3)}$ $(\bar{\alpha} = 4, \ldots, D)$ of the equations of state in the extra dimensions.

First, we defined the gravitational field (i.e. the metric coefficients) of the many-body system, since we need it to build the Lagrange function. We considered the weak field limit, assuming that the gravitational field is weak and velocities of masses are much less than the speed of light. Then, we used the found metric coefficients to construct the Lagrange function of the system. We proved that the Lagrange function exists only if $\Sigma \equiv \sum_{\bar{\alpha}} \omega_{(\bar{\alpha}-3)} = 0$. This strongly contradicts the observations, e.g., the PPN parameter γ, which demand $\Sigma \approx (3 - D)/2 \sim O(1)$ for $D > 3$. Hence, we come into conflict.

It is not difficult to realize that this result can be generalized to the case of arbitrary Ricci-flat internal spaces. Calabi-Yau manifolds are of special interest because these Ricci-flat spaces are widely used in superstring theories. It is also of interest to investigate the many-body problem in the case of KK models with spherical compactification of the internal spaces considered, e.g., in our papers [12, 13]. We will elaborate both of these cases in our forthcoming papers.

One more important result obtained in this paper is worth noting. It is well known [5–7] that tension in the internal spaces is the necessary condition to satisfy the gravitational experiments in KK models with toroidal compactification. In our paper, we have shown that the presence of pressure/tension in the internal space leads necessarily to the uniform smearing of the gravitating masses over the internal space. For example, black strings/branes have tension in the internal space (see, e.g., [14]). Therefore, they should be smeared. However, uniformly smeared gravitating bodies cannot have excited KK states (KK particles), which looks unnatural from the point of quantum mechanics.

To conclude our paper we state on the grounds of our investigations that considered multidimensional Kaluza-Klein models face severe problems.

ACKNOWLEDGEMENTS

This work was supported in part by the "Cosmomicrophysics-2" programme of the Physics and Astronomy Division of the National Academy of Sciences of Ukraine.

References

[1] Th. Kaluza, *Zum Unitätsproblem der Physik*, Sitzungsber. d. Preuss. Akad. d. Wiss., 966 (1921); O. Klein, *Quantentheorie und fünfdimensionale Relativitätstheorie*, Zeitschrift für Physik **37**, 895 (1926).

[2] J. Polchinski, *String Theory, Volume 2: Superstring Theory and Beyond* (Cambridge University Press, Cambridge, 1998).

[3] G. Bhattacharyya, A. Datta, S.K. Majee and A. Raychaudhuri, Nucl. Phys. B **821**, 48 (2009); arXiv:hep-ph/0904.0937.

[4] M. Eingorn and A. Zhuk, Class. Quant. Grav. **27**, 205014 (2010); arXiv:gr-qc/1003.5690.

[5] M. Eingorn and A. Zhuk, Phys. Rev. D **83**, 044005 (2011); arXiv:qr-qc/1010.5740.

[6] M. Eingorn, O. de Medeiros, L. Crispino and A. Zhuk, Phys. Rev. D **84**, 024031 (2011); arXiv:gr-qc/1101.3910.

[7] M. Eingorn and A. Zhuk, Phys. Lett. B **713**, 154-159 (2012); arXiv:gr-qc/1201.1756.

[8] L.D. Landau and E.M. Lifshitz, *The Classical Theory of Fields, Fourth Edition: Volume 2 (Course of Theoretical Physics Series)* (Pergamon Press, Oxford, 2000).

[9] C.M. Will, *Was Einstein Right? Testing Relativity at the Centenary.* In *100 Years of Relativity: Space-time Structure: Einstein and Beyond*, edited by A. Ashtekar (World Scientific, Singapore, 2005); arXiv:gr-qc/0504086.

[10] Bh. Jain and J. Khoury, *Cosmological Tests of Gravity* (2010); arXiv:astro-ph/1004.3294.

[11] B. Bertotti, L. Iess, and P. Tortora, Nature **425**, 374 (2003).

[12] A. Chopovsky, M. Eingorn and A. Zhuk A, Phys. Rev. D **85**, 064028 (2012); arXiv:qr-qc/1107.3388.

[13] A. Chopovsky, M. Eingorn and A. Zhuk A, Phys. Rev. D **86**, 024025 (2012); arXiv:qr-qc/1202.2677.

[14] D. Kastor and J. Traschen, JHEP **0609**, 022 (2006); arXiv:hep-th/0607051.